

Intracellular selection of peptide inhibitors that target disulphide-bridged A β ₄₂ oligomers

Nicola Acerra, Neil. M. Kad, Harish Cheruvara, and Jody M. Mason*

School of Biological Sciences, University of Essex, Wivenhoe Park, Colchester, Essex CO4 3SQ, United Kingdom

Received 1 May 2014; Accepted 18 June 2014

DOI: 10.1002/pro.2509

Published online 20 June 2014 proteinscience.org

Abstract: The β -amyloid (A β) peptide aggregates into a number of soluble and insoluble forms, with soluble oligomers thought to be the primary factor implicated in Alzheimer's disease pathology. As a result, a wide range of potential aggregation inhibitors have been developed. However, in addition to problems with solubility and protease susceptibility, many have inadvertently raised the concentration of these soluble neurotoxic species. Sandberg *et al.* previously reported a β -hairpin stabilized variant of A β ₄₂ that results from an intramolecular disulphide bridge (A21C/A31C; A β _{42cc}), which generates highly toxic oligomeric species incapable of converting into mature fibrils. Using an intracellular protein-fragment complementation (PCA) approach, we have screened peptide libraries using *E. coli* that harbor an oxidizing environment to permit cytoplasmic disulphide bond formation. Peptides designed to target either the first or second β -strand have been demonstrated to bind to A β _{42cc}, lower amyloid cytotoxicity, and confer bacterial cell survival. Peptides have consequently been tested using wild-type A β ₄₂ via ThT binding assays, circular dichroism, MTT cytotoxicity assays, fluorescence microscopy, and atomic force microscopy. Results demonstrate that amyloid-PCA selected peptides function by both removing amyloid oligomers as well as inhibiting their formation. These data further support the use of semirational design combined with intracellular PCA methodology to develop A β antagonists as candidates for modification into drugs capable of slowing or even preventing the onset of AD.

Keywords: amyloid; protein misfolding; protein-protein interactions; library screening; protein-fragment complementation assay; Alzheimer's disease

Introduction

It has been widely speculated that small soluble oligomers of A β play a major role in the pathology of

Alzheimer's disease (AD).^{1–4} However, developing drugs that prevent the accumulation of these species has proven difficult. Initial studies that sought to

Abbreviations: A β ₄₂, β -amyloid 1–42 variant; CD, circular dichroism; HFIP, hexafluoroisopropanol; MTT, (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; PCA, protein-fragment complementation assay; PPI, protein-protein interaction; TEM, transmission electron microscopy; TFA, trifluoroacetic acid; ThT, Thioflavin-T; OAF, oblique angle fluorescence; AFM, atomic force microscopy

Additional Supporting Information may be found in the online version of this article.

Grant sponsor: Pilot project (Alzheimer's Research UK; J.M.M.); Grant number: ART/PPG2008B/2. Grant sponsor: AgeUK [New Investigator Award (#304)]. Grant sponsor: Cancer Research UK Career Establishment Award (J.M.M.); Grant number: C29788/A11738. Grant sponsor: Wellcome Trust Project; Grant number: WT090184MA. Grant sponsor: BBSRC (New Investigator Award N.M.K.); Grant number: BB/1003460/1. Grant sponsor: University of Essex Departmental Studentship (N.A.). Grant sponsor: Parkinson's UK PhD Studentship (H-1001; J.M.M and N.M.K).

*Correspondence to: Jody M. Mason; School of Biological Sciences, University of Essex, Wivenhoe Park, Colchester, CO4 3SQ, United Kingdom. E-mail: jmason@essex.ac.uk

breakdown large insoluble oligomers have in some instances led to an accumulation of small toxic species,⁵ while more recent work has shown that even dimers or trimers of A β are toxic and able to impair synaptic function.⁶ In screening libraries to identify peptides capable of modifying aggregation, an elegant approach would be to lock A β_{42} into a toxic conformation. This would ensure population of an oligomeric state that is pathogenically relevant to the disease. Sandberg *et al.* previously studied the effect of an intramolecular disulphide-bridged double mutant of A β_{42} (A21C/A31C; A β_{42cc}).⁷ In this approach, conservative mutations were used to create a disulphide bridge and force an otherwise unstructured and monomeric A β_{42} into the β -hairpin conformation found in mature fibrils. The conformation, therefore, provides the structural basis for subsequent amyloidogenesis.⁸ Disulphide tethering was found to stabilize highly toxic A β oligomers that retained the conformational and biological properties of wild-type A β . Moreover, these molecules did not go on to convert into mature fibrils and were shown to contain SDS-stable dimeric and trimeric species. Experiments using SH-SY5Y human neuroblastoma cells demonstrated that A β_{42cc} oligomers or protofibrillar species formed by these oligomers were 50 times more potent inducers of neuronal apoptosis than amyloid fibrils or samples of monomeric wild-type A β_{42} , in which toxic aggregates were only transiently formed.

A wide variety of approaches have been used to create amyloid inhibitors, including small molecule, antibody, and peptide-based inhibitors. These have ranged from those that stabilize the native state, sequester monomers, inhibit amyloid growth, as well as those that initiate clearance via chaperones and proteases (see Ref. 9 for an overview). In our own studies, we previously presented a novel variation of the intracellular protein-fragment complementation assay (PCA) selection system whereby bacteria were used to screen peptide libraries and generate A β aggregation antagonists¹⁰ and subsequently their retroinverted analogues.¹¹ During PCA a “core recognition element” A β_{25-35} , was used as the target since, along with A β_{15-21} , it is known to form amyloid fibrils in isolation and instigate amyloid formation in the parental protein. A β_{25-35} was fused to one half of murine dihydrofolate reductase (mDHFR) with a peptide library fused to the second half. Library members that bound A β_{25-35} resulted in a colony under selective conditions. The most effective binders were then further enriched by growth competition to isolate from binders that led to lower growth rates. Reduced bacterial growth rates are due to the inherent propensity for A β to aggregate and lead to increased cytotoxic effects.^{12,13} Three outcomes are possible for any given library member:

- A. Library members bind A β , reduce its toxicity and recombine mDHFR to confer cell survival.
- B. Library members bind A β and recombine mDHFR but either populate or do not prevent population of a toxic species. These result in reduced cell growth relative to 1, or cell death.
- C. Library members with no affinity for A β and, therefore, no effect on amyloid formation will not recombine mDHFR, resulting in cell death.

Intracellular selection means no assumptions are made regarding the mechanism of antagonist action or which amyloid state becomes populated during selection, aiding the removal of molecules with undesirable properties from the library such as those that are too hydrophobic hence insoluble, those tagged for degradation, those susceptible to protease action, and those that are nonspecific.

To further strengthen this approach, we report on the use of full length A β_{42cc} as the target molecule rather than A β_{15-21} or the A β_{25-35} amyloidogenic regions of the molecule used previously.¹⁰ In this experiment, PCA is undertaken using A β_{42cc} under oxidizing conditions to ensure that:

1. During selection the A β_{42cc} target is constrained into the β -hairpin structure associated with the pathology of the disease
2. The “core recognition elements” of A β_{42cc} associated with instigation of amyloid formation remain exposed, largely unmodified, and in the correct conformation.
3. A β_{42cc} is able to form soluble oligomeric structures that are highly toxic and associated with AD pathology, thus ensuring that a larger number of structurally relevant epitopes are presented relative to larger and/or insoluble fibrils populated in the absence of the bridge.
4. Expressing the increased toxicity variant A β_{42cc} raises the sensitivity of the PCA relative to A β_{42} . This is because cells in scenario B above are predicted to die more readily or grow at reduced rates.

We find the peptides derived using this approach to be capable of binding to the disease relevant wild-type A β_{42} and reducing associated amyloid formation. To understand the mechanism of peptide action we have studied the effectiveness of peptides and found them to be capable of removing preformed fibrils as well as preventing amyloid from forming. These studies have successfully produced a number of lead peptide sequences that are expected to provide a scaffold for future drug candidates.

Results

We have combined semirational design with intracellular selection using amyloid-PCA to screen peptide




Figure 1. The PCA for amyloid systems. Selection is undertaken using SHuffle cells that harbor an oxidizing cytoplasmic environment to promote disulphide bond formation. Library members that bind to $A\beta_{42cc}$ lead to the recombination of murine DHFR and lead to a colony under selective conditions (bacterial DHFR is specifically inhibited using trimethoprim). Subsequent competition selection in liquid media isolates winners of highest efficacy. Those library members that bind the target and are able to confer faster growth rates by reducing the toxic effects of the amyloid protein most effectively will be selected. As the assay is performed in the cytoplasm of *E.coli*, any nonspecific, unstable, aggregation prone (insoluble), protease susceptible members are removed.

libraries and select $A\beta_{42}$ aggregation antagonists.^{10,14,15} A disulphide-bridged variant of $A\beta_{42}$, known as $A\beta_{42cc}$, has been used as the target for library screening since it constrains the protein into the β -hairpin structure found in amyloid structures that form oligomers associated with AD toxicity (Fig. 1). PCA libraries were initially screened and selected on M9 minimal agar plates. Following this initial single-step selection, colonies underwent competition selection where they were pooled and grown before dilution into liquid M9 minimal media. This process was repeated multiple times as passages to select for the most effective binding sequence. In this process, 1–2 high affinity binders that inhibit amyloid formation and reduce bacterial toxicity were found to display the fastest growth and dominate the bacterial pool. The selection process indicated that only a very limited subset of library members were able to bind $A\beta_{42cc}$ (<1%) and that these can be further separated during competition selection PCA to isolate the most effective binders. DNA sequencing continued throughout passaging until selection arrived at one discrete peptide. All subsequent aggregation studies were undertaken using $A\beta_{42}$ as $A\beta_{42cc}$ is not found naturally, with ThT binding and TEM data demonstrating that fibrils do not form when the intramolecular disulphide bridge is intact.⁷ We demonstrate that the sequences identified using the $A\beta_{42cc}$ target can be transferred to the biologically relevant $A\beta_{42}$.

Library generation

PCA was undertaken using an $A\beta_{42cc}$ target with sequences corresponding to $A\beta_{15-21}$ (QKLVFFA) $A\beta_{29-35}$ (GAIIGLM), and $A\beta_{36-42}$ (VGGVVIA) as library templates. The first of these two regions are known to aggregate into toxic fibrils in isolation¹⁶⁻¹⁸ and therefore provided a strong starting point for deriving peptides known to bind to $A\beta_{42cc}$ and potentially capable of inhibiting aggregation. The third sequence incorporates the two additional C-terminal amino acids (Ile-Ala) that renders $A\beta_{42}$ much more toxic than $A\beta_{40}$. The first library incorporated four fully randomized residues at positions 17–20 of $A\beta_{15-21}$, generating a library size of 160,000 (¹⁵QKxxxxA²¹), and resulted in one winner (cys₁₅₂₁; Table I). Interestingly, this library had previously resulted in no colonies when using either $A\beta_{42}$ or $A\beta_{15-21}$ as target, suggesting an improved epitope for binding on introduction of the cysteine-bridge. Moreover, the selected sequence, QKVLLFA, bore remarkable similarity to the $A\beta_{15-21}$ (QKLVFFA) parental sequence on which it was based, having been enriched during PCA from four fully randomized residues at positions 17–20. The $A\beta_{15-21}$ peptide has been previously shown to form amyloid in isolation and found to be toxic in MTT experiments.^{19,20} The second library incorporated four fully randomized positions at 29–30 and 34–35 of $A\beta_{29-35}$, with residues 31–33 fixed from a previous selection,¹⁰ generating a 160,000 member library (²⁹xxKATxx³⁵),

Table I. PCA Derived Sequences

Name	Sequence
cys ₁₅₂₁	AC-QK <u>VLLFA</u> -NH ₂
cys _{2935a}	AC- <u>AGKATGL</u> -NH ₂
cys _{2935b}	AC- <u>GAKATAN</u> -NH ₂
cys ₃₆₄₂	AC- <u>RWGVVWG</u> -NH ₂
iAβ5	AC-LPFFD-NH ₂

Four libraries consisting of 160,000 members each and based around residues 15–21, 29–35, and 36–42 of the Aβ_{1–42} were fully randomised at positions shown underlined to allow options for all amino acids. A second library consisting of 160,000 members was fully randomised at positions 29–30 and 34–35 with the KAT sequence from the first selection fixed. Cys_{2935a/b} is based on a previously selected sequence that targeted Aβ_{29–35}.¹⁰ Acetylation and amidation of the protein was added to mimic the polypeptide backbone and provide additional potential hydrogen-bond acceptors and donors. The positive control sequence, iAβ5, is also shown.

and resulted in two amyloid-PCA winners (cys_{2935a} and cys_{2935b}; Table I) with no sequence homology to the amyloid template on which they were based. The last library fixed residues 38–40 of Aβ_{36–42} as “GVV” as is found in wild-type Aβ₄₀ and Aβ₄₂, but was fully randomized at residues 36–37 and importantly at residues 41–42 associated with the increased toxicity of Aβ₄₂, generating a 160,000 member library (³⁶xxGVVxx⁴²), and resulting in one winner (cys₃₆₄₂; Table I).

Peptide characterization

PCA derived peptide sequences (Table I) were synthesized and characterized using a number of experiments that included thioflavin-T (ThT) dye binding, circular dichroism (CD), oblique angle fluorescence (OAF), and atomic force microscopy (AFM). The results demonstrate that the peptides do not aggregate in isolation, and are able to prevent Aβ₄₂ aggregation and/or remove preformed fibrils. In addition, growth competition assays using *E.coli* under PCA conditions in M9 media and an MTT assay using PC12 cells, both using the Aβ₄₂ parent peptide, were undertaken to establish cytotoxicity to bacterial and mammalian cells. The growth competition experiments simultaneously demonstrate that peptides bind to Aβ₄₂ and reduce its associated toxicity during bacterial selection. MTT experiments were used to establish that the toxicity associated with extracellular Aβ₄₂ to mammalian cells could be reduced when incubated in the presence of PCA selected peptides.

Cell growth experiments

The effect of inhibitors on the growth of *E.coli* harboring pES300d-Aβ_{42cc}-DHFR2 target and pES230d-antagonist-DHFR1 fusion plasmids as present in the final PCA selection round were tested (Fig. 2). In this experiment, cells were grown in a shaking incubator from a starting OD₆₀₀ of 0.02 under PCA

conditions in M9 minimal media containing Cm, Amp, and Kan to retain target and antagonist expressing plasmids as well as pREP4 for expression of the lac repressor. In addition Tmp was included for inhibition of bacterial DHFR and IPTG to induce high levels of target and antagonist expression. This experiment monitors both mDHFR reassembly, and, therefore, binding of antagonist to the Aβ_{42cc} target, as well as the toxicity of the oligomeric state that is populated. As expected, expression of the toxic Aβ_{42cc} did not result in significant levels of growth even though the protein is well documented to self-associate (Fig. 2). In addition, western blots (Supporting Information Fig. S5) show that Aβ_{42cc} is expressed in the soluble fraction, suggesting that the protein is both soluble and toxic and is, therefore, populating toxic protofibrillar structures. All four antagonists in this study, along with the positive control cJun-FosW, were clearly able to restore bacterial growth thereby providing strong evidence for direct binding and reduced toxicity in the context of this bacterial selection system.

ThT experiments demonstrate a reduction in amyloid content

To determine the ability of PCA derived peptides to prevent fibril assembly (inhibition) or to breakdown preformed fibrils (reversal), ThT fluorescence was used as an indicator of the degree to which Aβ₄₂ aggregates into amyloid fibrils. In these assays, Aβ₄₂




Figure 2. To confirm that expression of Aβ_{42cc}-DHFR1 / Aβ_{42cc}-DHFR2 fusions impedes the growth rate of *E. coli*, and to ascertain substitution with antagonist-DHFR1 fusions are able to reverse this effect, growth competition experiments were undertaken in M9 liquid media in an identical manner to that during the PCA selection process. In these experiments, cells expressed either i) Aβ_{42cc} + Aβ_{42cc}, ii) Aβ_{42cc} + antagonist, or iii) nontoxic cJun + FosW. In this experiment, only scenarios ii) and iii) led to significant growth rates in *E. coli*, providing evidence that a high affinity interaction is formed and that toxicity associated with expression of Aβ_{42cc} has been lowered, leading to significant growth rates relative to i).




Figure 3. ThT Inhibition and Reversal data for a) individual peptides and b) peptides used in combination. The data show the effects of different stoichiometries of the inhibitors cys₁₅₂₁, cys_{2935a}, cys_{2935b}, cys₃₆₄₂, cys₁₅₂₁ + cys_{2935a}, cys₁₅₂₁ + cys₃₆₄₂, and iAβ₅ on the aggregation of 50 μM Aβ₄₂. The assay was undertaken at 3 days for the inhibition assay. For the reversal assay, Aβ₄₂ was incubated in isolation before addition of peptides on Day 3 with the assay undertaken on Day 6. The assay was performed at 10 μM Aβ₄₂ concentrations. All errors are expressed as the standard error of the mean.

was rendered monomeric²¹ and freeze-dried before being resuspended and incubated at 37°C, leading to formation of amyloid aggregates. For the inhibition assay, PCA-selected peptides were added on Day 0 and their ability to prevent amyloid formation was assessed after 3 days. For the reversal assay Aβ₄₂ was incubated in isolation for three days to allow amyloid formation before addition of peptide, followed by incubation for a further 3 days to monitor the ability of peptides to breakdown amyloid aggregates. In all cases, a positive control peptide from the literature (iAβ₅^{22,23}) known to perform well in ThT assays was used. Figure 3 shows the results of inhibition and reversal experiments for the four

single PCA-selected peptides and two examples of peptides administered in combination. Both inhibition and reversal experiments have been undertaken at five different Aβ₄₂:peptide ratios ranging from 10-fold substoichiometric to 10-fold superstoichiometric. For every peptide, and at the majority of stoichiometries, the ThT-bound signal was reduced for both inhibition and reversal experiments, indicating that peptides are able to bind Aβ₄₂ and lower the amount of ThT-bound amyloid in solution. Consistent with previous studies, changes in Aβ₄₂:inhibitor stoichiometry altered binding although it was not clear if this was dependent on dose.^{10,24} However, on average the effects observed at ratios of 1:0.1 are small

while ratios of 1:1 are lower than 1:2 or 1:4 (see Supporting Information Fig. S4). These ratios were required to consistently elicit an effect in both inhibition and reversal experiments. Increasing the ratio to 1:10 led to poorer levels of inhibition for several peptides (and consequently increased ThT fluorescence). What is also clear from these experiments is that ThT-bound amyloid can be frequently reduced to less than 50% of the signal found in the A β ₄₂ positive control, with peptides performing well in both inhibition and reversal experiments. Reassuringly, the positive control iA β 5 peptide was also able to reduce the ThT signal by >40%.

In addition, we have conducted experiments where amyloid-PCA derived peptides predicted to target different regions of the disulphide tethered A β _{42cc} have been combined [Supporting Information Fig. 3(b)]. These combinations were (a) cys₁₅₂₁/cys_{2935a} and (b) cys₁₅₂₁/cys₃₆₄₂. In both instances, these peptides were derived from libraries based on (and, therefore, predicted to independently target) the first and second strands within the tethered β -hairpin structure of A β _{42cc}. Indeed, cys_{2935a} is based on a sequence that was selected using the A β ₂₅₋₃₅ target,¹⁰ with the same library unable to yield any binding sequences when using the wild-type A β ₄₂ or A β ₁₅₋₂₁ as a target. In addition, the library used to generate cys₁₅₂₁ was also unable to generate any hits against wild-type A β ₄₂. This is significant since it suggests that A β _{42cc} represents a far more accessible target for intracellular library screening and antagonist selection. Despite this, ThT experiments in which these peptides were combined generated only one instance where two peptides assayed together were, within error, consistently more effective across the stoichiometries than the average of the individual component peptides [Fig. 3(b)]. This was observed for inhibition experiments that combined cys₁₅₂₁ and cys_{2935a}. In this case, an additional 25–35% reduction in ThT fluorescence was observed over the average of the individual peptides at 1:0.1, 1:2, 1:4, and 1:10 stoichiometries, indicating a benefit in combining them [Fig. 3(b), green vs. purple]. At 1:1, large cumulative errors precluded any interpretation. In contrast, combinations of cys₁₅₂₁ and cys₃₆₄₂ gave no consistent benefit in inhibition experiments. For peptides tested in combination during reversal experiments, no consistent benefit was observed over either peptide assayed alone. Rather, combinations of cys₁₅₂₁ and cys_{2935a} displayed signs of synergy at 1:1 to 1:2 molar ratios with the trend reversing for 1:4 and 1:10, indicating that higher concentrations of peptide can lead to increased ThT fluorescence. As increasing the molar ratio did not lead to improvements in the reversal of amyloid in most cases, this indicated that a two-fold excess of peptide:A β may be sufficient to reverse fibril formation. A similar trend was true of combining cys₁₅₂₁ and cys₃₆₄₂ in reversal experiments. However, in both cases the trend was not observed in inhibition

assays, perhaps reflecting the fact that more binding sites are available in monomerized A β ₄₂ to which inhibitor can bind and, therefore, remain soluble relative to aged and aggregated samples. Lastly, it cannot be ruled out that the combined peptides do not interact with each other; however, mixing peptides does not result in any gain in amyloid formation as increased ThT signals are not observed.

CD experiments demonstrate a reduction in β -sheet content

As amyloid fibrils are predominately β -sheet, we used CD spectroscopy to provide an end-point characterization of the aggregates formed once ThT experiments were complete. Consistent with a β -sheet structure, spectra show a negative peak at 218 nm. At higher molar ratios (1:2 or greater for inhibition and 1:4 or greater for reversal) an additional large minima of up to -11 millidegrees was observed at 200 nm for the majority of peptides, indicating the adoption of a mixture of β -sheet (A β ₄₂) and random coil-like structures (peptides). This minima was noticeably less pronounced for the cys₃₆₄₂ peptide and was not observed for iA β 5, which adopted a spectrum consistent with a β -sheet structure at all concentrations. During CD experiments, we observed a decrease in β -sheet content in the majority of samples, supporting the ThT data by demonstrating that peptides reduce the global β -sheet content relative to the A β ₄₂ sample and, therefore, the amyloid content. These experiments therefore suggest that peptides can bind A β ₄₂ and exert their effect by reducing the amount of protein in the amyloid form. Lastly, ThT and CD spectroscopy experiments undertaken on peptides in isolation that have been incubated at 50 μ M for 3 days under conditions identical to those used in aggregation assays using A β ₄₂ demonstrate that peptides do not bind significant amounts of ThT, and that the CD signal for all peptides (at 0:1) is consistent with that of a random coil or weakly helical conformation.

OAF microscopy

Samples used in ThT and CD experiments were also imaged using OAF microscopy for both inhibition and reversal experiments. To prevent bias toward any one sample the experiment was carried out blind.²⁵ This technique allows for surface associated and stacked aggregates of amyloid fibers to be directly imaged. It was possible to assess the amount of protein deposited as amyloid and its morphology. OAF experiments were undertaken for both inhibition and reversal at all of the molar ratios assayed in ThT experiments (Fig. 4). In almost every case, for both inhibition and reversal experiments, a reduction in the amount of amyloid was observed relative to A β ₄₂ incubated in isolation under




Figure 4. OAF Microscopy data for all inhibitors at all stoichiometries. During inhibition experiments, Aβ₄₂ was grown with inhibitor for 3 days and assayed for peptide induced inhibition of amyloid formation. During reversal experiments, Aβ₄₂ was grown without inhibitor for 3 days, after which inhibitor was added at a stoichiometry of 1:1 and followed by a further 3 days incubation to assay for peptide induced reversal of amyloid deposition. Each sample was then imaged by fluorescence microscopy and panels showing representative images obtained. To quantify amyloid deposition the mean grey value over a 160 px × 160 px (13 μm × 13 μm) area randomly chosen for five separate images is plotted as fluorescence intensity. Each data point is scaled to overcome the “background noise” by taking Aβ (1:0) as the maximum and iAβ5 as the minimum [i.e., (signal-iAβ5)/(Aβ₄₂-iAβ5)]. This defines the range over the positive and negative controls.

identical conditions. In a number of cases amyloid deposits much smaller in size were observed.

Atomic force microscopy

Again samples used in ThT and CD experiments were imaged using AFM as a second direct

qualitative measure of fibril formation (Fig. 5). A stoichiometry of 1:2 (50 μM:100 μM) was chosen for AFM experiments as this was the concentration at which peptide inhibitors were on average deemed to be most effective in ThT experiments. AFM was undertaken for both inhibition and reversal experiments on




Figure 5. AFM data. To demonstrate that inhibitors cys_{1521} , cys_{2935a} , and $cys_{2935a} + cys_{1521}$ do not aggregate in isolation, they were grown alone for 3 days at a concentration of $100 \mu M$ (i.e., 0:2 molar ratio) under the same conditions as inhibition experiments. In all cases, no fibril formation was observed. Inhibition experiments were undertaken at an $A\beta_{42}$:inhibitor stoichiometry of 1:2 (row two). In this experiment, the same sample was used as for ThT experiments, in which $A\beta_{42}$ was grown with inhibitor for 3 days at $37^\circ C$. All of the peptides are inhibitory, in particular the combination of $cys_{2935} + cys_{1521}$. For reversal experiments, $A\beta_{42}$ was grown alone for 3 days, after which inhibitor was added at a stoichiometry of 1:2 and followed by a further 3 days incubation to assay for peptide induced reversal of amyloid deposition. Each of the samples were imaged by AFM and panels showing representative images obtained. Also in reversal experiments, it can be clearly observed that for all of inhibitors there is a reduction of fibril, although this is less pronounced than for inhibition experiments. Scale bars are shown.

cys_{1521} , cys_{2935a} , and $cys_{1521} + cys_{2935a}$ as an additional measure of the extent to which these peptides are effective when mixed. A major reduction in the amount of amyloid was observed for all samples relative to the $A\beta_{42}$ control. No fibrils were observed for peptides in the absence of $A\beta_{42}$, either alone or when mixed. For inhibition experiments in the case of $cys_{1521} + cys_{2935a}$ no fibrils were observed; with mixing appearing to remove fibrils present for the cys_{1521} sample. For reversal experiments, the residual fibrils observed in the cys_{2935a} were also largely removed on mixing with cys_{1521} .

Cell-toxicity experiments

MTT ((3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide)) cell toxicity experiments were performed using Rat pheochromocytoma (PC12) neuronal-like cells to reflect the nature of toxicity in the disease via the reduction of a redox dye (Fig. 6). We assessed the toxicity of extracellular $A\beta_{42}$ deposits on PC12 cell integrity and its amelioration by preincubation of $A\beta_{42}$ with peptides. Cells incubated with buffer alone resulted in a reduction of the redox dye MTT, leading to a color change from

yellow to purple that can be measured by a change at A_{540} . In contrast, incubation of cells with $A\beta_{42}$ resulted in a large decrease in cell viability and consequently the ability to reduce MTT, leading to a smaller signal change. To study, the ability of inhibitors to reverse the effect on cell viability MTT assays were performed with a range of molar ratios relative to $A\beta_{42}$ and normalized relative to cells in isolation (normalized as 0% death) and cells incubated with $A\beta_{42}$ alone (normalized as 100% death). Although the changes are small and the cumulative errors resulting from comparatively small changes can be large, we were nonetheless able to draw some general conclusions from this assay. In particular, we observed that effective peptides were able to generate modest reductions in toxicity of 5–20% in the majority of cases. Neither cys_{2935a} or cys_{2935b} appeared effective in this assay, displaying almost no improvement across all stoichiometries. Cys_{1521} was able to elicit a small decrease of 5–15% in a nondose dependent manner. Cys_{3642} demonstrated a decrease across all molar ratios of 10–20%. A combination of $cys_{1521} + cys_{2935a}$ or $cys_{1521} + cys_{3642}$ did not conclusively lead to additional lowering of




Figure 6. Shown are a) the results from the MTT toxicity assay using Aβ₄₂ and selected inhibitors using different molar ratios after 24 h of incubation with PC12 cells. The assay was performed with 10 μM Aβ₄₂ and different concentrations of inhibitor, for example, 1:0.1 (1 μM), 1:1 (10 μM), 1:4 (40 μM), 1:10 (100 μM). b) a comparison of the MTT results for individual peptides against those used in combination. All errors are expressed as the standard error of the mean.

toxicity [Fig. 6(b)]. Although reductions in toxicity greater than the average of the component peptides were observed, these were small; 5% for cys₁₅₂₁/cys_{2935a} at 1:10 and 10% for cys₁₅₂₁/cys₃₆₄₂ at 1:10. At other molar ratios the errors were either too large or no additional benefits over Aβ₄₂ alone were observed. It should be noted that as PCA is undertaken inside bacteria it is possible that intracellular selection of peptides that are nontoxic to *E. coli* may not be as effective when transferred into a mammalian cell system. There, exposure to different off-target interactions and a different protease pool will occur, in addition to the necessity for the peptide to function in the extracellular space, where Aβ is known to aggregate in the brain. This may explain the discrepancy between the effects on growth rates in *E. coli* and efficacy in MTT experiments on peptides incubated with PC12 cells.

Discussion

We have used an intracellular PCA combined with semirational library design to identify peptides that can reverse and inhibit Aβ₄₂ aggregation by up to 80% as well as lead to a modest reduction (5–20%) in its

toxicity. As a target during bacterial selection, we utilized a double cysteine mutant Aβ_{42cc} (A21C/A30C⁷) that has been shown to lead to the population of pathogenically relevant soluble oligomers in the cytoplasm of an *E. coli* strain that harbors an oxidizing cytoplasm environment.^{10,26} An advantage of using PCA with Aβ_{42cc} is that the disulphide bridge forces Aβ into the β-hairpin conformation that is associated with toxicity in AD. The Aβ_{42cc} mutant has been shown to lead to the population of highly toxic oligomers, suggesting that this mutant can be used to improve the stringency of the amyloid-PCA assay. In addition, as oligomers are adopted, we envisage that this conformationally restricted mutant generates an increase in the number of exposed structural epitopes relative to fibrils formed by wild-type Aβ₄₂, meaning that more mDHFR activity can be attained within cells harboring binders of this disulphide mutant. To target, the two β-strands either side of the hairpin, we used Aβ_{15–21}, Aβ_{29–35}, and Aβ_{36–42} as design scaffolds and fully randomized residues at four positions to create three libraries of 160,000 members. All three libraries resulted in the identification of sequences able to bind the target sequence. In the case of wild-type Aβ₄₂, it

should be noted that it is possible that the peptide inhibitors may be able to function by stabilizing monomeric A β ₄₂ before it misfolds into the β -hairpin structure, thus preventing conversion into a pathogenic structure. By using the restrained A β _{42cc} protein, and because we were unable to identify full length A β ₄₂ binders using many of the same libraries, we infer that the wild-type structure is a poor target for antagonist recognition, possibly due to a lack of well-defined structural epitopes in the nonamyloid or early oligomeric conformations of the molecule. Thus, the generalized approach of using A β _{42cc} to screen libraries and generate peptide hits before testing selected peptides on wild-type A β ₄₂ is potentially very exciting.

Combination experiments

As previously noted, we have identified hits from the same 15–21 library that was unable to yield binders for either wild-type A β ₄₂ or A β _{25–35} target proteins (unpublished data). The failure to identify binders in these previous PCA screens suggests that the epitope presented was insufficient for effective binding. Thus, we believe that the restricted conformation of A β ₄₂ in the disulphide bridged version represents an improved target for our “recognition-element” based libraries. As the 15–21 library was unable to provide colonies under PCA conditions with A β ₄₂, we instead focused our efforts on the 25–35 region of the molecule. We therefore envisage that the cys₁₅₂₁ winner is able to target the first β -strand in the constrained amyloid conformation of A β _{42cc}, with the cys₂₉₃₅ and cys₃₆₄₂ libraries targeting the second β -strand. To look for synergy when used in combination, we therefore incubated the cys₁₅₂₁ winner with either cys_{2935a} or cys₃₆₄₂. In the case of cys₁₅₂₁ and cys_{2935a}, we found a reduction in bound-ThT signal of 20% over the average of the two individual peptides, indicating that there is an added benefit in combining peptides, and that they may occupy different regions within A β . MTT data produced large errors, however, improvements in cell viability of 5–10% at some molar ratios were observed, indicating that the combined effect of the inhibitors is greater than the average of the two. In agreement with this, OAF and AFM experiments also suggest that there is an additional benefit in mixing the cys₁₅₂₁ and cys_{2935a} peptides.

This study has led to a number of peptides that are capable of preventing assembly of amyloid aggregates as well as breaking down preformed amyloid and further investigation is warranted in the search for molecules that can ultimately slow or even prevent the onset of AD.

Materials and Methods

Mutagenesis and protein engineering

pES300d-A β _{42cc}-DHFR2 was created using overlap extension PCR with codon changes corresponding to

A21C and A30C. These sites have been previously chosen to constrain the peptide into a β -hairpin conformation⁷ and are located on nonhydrogen bonded sites between β -strands either side of the hairpin and in close proximity to each other.²⁷

Choice of *E. coli* strain

PCA were undertaken using *E. coli* SHuffle Express cells (New England Biolabs). SHuffle cells have been engineered to possess an oxidative cytoplasmic environment that favors disulphide bond formation. In particular, SHuffle cells have been engineered to lack thioredoxin reductase (*trxB*) and glutathione reductase (*gor*) with an additional suppressor mutation (*ahpC*) which is required to restore viability, allowing the formation of stable disulphide bonds in the cytoplasm. Under these conditions, thioredoxins are in their oxidized state, converting them from reductases to oxidases. Proteins that require disulphide bonds for their folding can, therefore, be oxidized and form stable disulphide bonds within the cytoplasm. This cell line also constitutively expresses a chromosomal copy of the disulphide bond isomerase DsbC. DsbC promotes the correction of misoxidized proteins into their correct form.^{10,26}

Single step selection PCA

pES300d-A β _{42cc}-DHFR2 and pREP4 (Qiagen; for expression of the lac repressor protein) were cotransformed into SHuffle Express cells (New England Biolabs) and plated onto LB agar with the appropriate antibiotics (Kan and Cm). These cells were next made electrocompetent before transformation with the appropriate pES230d-library-DHFR1 library plasmid (*E. coli* XL-1 cells were used for construction and cloning of libraries, as described previously^{10,14}). Transformed cells were next plated on three different media; 1/20th of the cells were plated onto LB agar with three antibiotics (Kan, Amp, and Cm) as a positive control of transformation efficiency. A further 1/20th of the solution was plated onto M9 minimal medium agar with 1 μ g/ml trimethoprim and the same three antibiotics as a negative control. Finally, the remaining 90% of transformed cells were plated onto M9 minimal agar in the presence of the three antibiotics, 1 μ g/ml trimethoprim, and 1 mM IPTG (isopropyl- β -D-thiogalactopyranoside), to induce expression of the two DHFR fragment fused peptides). This single-step PCA selection typically led to approximately 50–100 colonies from libraries of 160,000 members, meaning that at least 99% of all library members are removed at this stage owing to their inability to bind A β _{42cc} and reduce toxicity.

Competition selection PCA

To increase the selection stringency, growth competition experiments were undertaken. Using this

approach, selected colonies were pooled from the plate and grown in M9 minimal media under selective conditions (containing Kan, Amp, Cm, trimethoprim, and IPTG) and serially diluted over 5–10 passages. Using these sequential rounds of competition selection, subtle differences in growth rate are amplified, increasing the stringency of selection relative to the single-step method. Competition selection, therefore, allows the most effective 1–2 sequences to be isolated from the 50–100 A β_{42cc} binders that are initially identified during single step selection. At each passage glycerol stocks were prepared and sequencing results were obtained (GATC biotech, London) for DNA pools as well as individual colonies. For each passage, 50 μ L of liquid culture was added to 50 ml of fresh M9 minimal media, resulting in an approximate OD₆₀₀ of 0.01. Cells were incubated at 37°C until an OD₆₀₀ of \sim 0.4 was reached (typically 2–3 days), before moving to the next passage.

A β peptide preparation

A β_{42} was purchased as a pure recombinant peptide from rPeptide (Stratech) and was used for all of the experiments described. Prior to use, a quantity of peptide was weighed using an analytical balance before being treated to three rounds of dissolution in hexafluoro-2-propanol (HFIP), sonication, drying, dissolution in trifluoroacetic acid (TFA), sonication and drying, according to the Zagorski protocol,²¹ and then aliquoted into appropriately sized batches for subsequent assays and dried via lyophilisation before being dissolved in 10 mM potassium phosphate buffer (pH 7.4) to generate a final concentration of 50 μ M. TFA/HFIP treatment was used to ensure that amyloid growth always proceeds from the same monomeric state, thus reducing errors in amyloid formation measurements.

Peptide inhibitors

Peptides identified using amyloid-PCA (cys₁₅₂₁, cys_{2935a}, cys_{2935b}, cys₃₆₄₂, and a positive control from the literature, iA β_5 ²²), were obtained by Peptide Protein Research (Fareham, UK) as pure lyophilized peptides. Peptides were weighed using an analytical balance and stock solutions of 1 mM concentration were subsequently dissolved in ultrapure water. At this concentration (2–200x excess of that used in inhibition and reversal experiments) no aggregation or precipitate was observed. In addition, bioinformatics tools (e.g., Waltz,²⁸ Amylpred,²⁹ Pasta,³⁰ Zyggregator,³¹ and Tango³²) do not predict that any of the peptides contain amyloidogenic sequences or aggregate in isolation. Lastly, CD and ThT experiments on peptide antagonists in isolation demonstrate that these sequences do not bind ThT and form random coil-like species in isolation (see Supporting Information Fig. S1).

Growth competition experiments

To confirm that expression of A β_{42cc} -DHFR2 fragment fusions impedes the growth rate of *E. coli*, and to ascertain that amyloid-PCA derived peptides fused to the DHFR1 fragment are able to reverse this effect, growth competition experiments were undertaken. These experiments were performed in M9 minimal media in an identical manner to that during the PCA competition selection process. In these experiments (Fig. 2), cells expressed either (i) A β_{42cc} control, (ii) A β_{42cc} + PCA derived peptide, or (iii) the nontoxic control cJun+FosW. The latter PCA-derived pair is known to form a high affinity interaction, leading to significant growth rates relative to (i).

Circular Dichroism

Far-UV CD spectra were recorded on an Applied Photophysics Chirascan at 20°C using the same samples as for ThT experiments. Spectra were recorded over the 200–300 nm range at a scan rate of 10 nm/min with step size of 1 nm. Spectra were recorded as the average of two scans. Peptide (10 μ M in 10 mM Potassium Phosphate buffer pH 7.4) was added to a 0.1 cm cuvette (Hellma). Spectra were recorded as raw ellipticity.

Thioflavin T assays

ThT inhibition assays were performed with 50 μ M Zagorski treated²¹ monomeric A β_{42} in 200 μ L of 10 mM potassium phosphate, pH 7.4, with or without each peptide inhibitor at a concentration of 5 μ M (for 1:0.1 molar ratio), 50 μ M (for 1:1 molar ratio), 100 μ M (for 1:2 molar ratio), 200 μ M (for 1:4 molar ratio), and 0.5 mM (for 1:10 molar ratio). For the inhibition assays, the target-inhibitor mixtures were incubated together on day zero at 37°C. Single ThT readings were taken on day three at which maximal ThT binding was found. For the reversal assays, 200 μ L of 50 μ M target was incubated alone at 37°C for 3 days before adding to the required amount of lyophilized peptide. The vortexed target-inhibitor solutions were then incubated at 37°C for a further 3 days, during which time single ThT readings were taken on post-mix Day 3. Experiments were undertaken in duplicate and errors expressed as the standard error of the mean. The ThT assay solution was prepared from a 25 \times stock containing 500 μ M ThT in 250 mM Tris, pH 7.4. The stock was aliquoted and kept frozen until required. It was then allowed to thaw at room temperature for 10 min before dilution into the appropriate Tris buffer, giving the required freshly prepared ThT assay solution containing 20 μ M ThT in 10 mM Tris and buffer at pH 7.4. A total of 2960 μ L of the ThT assay solution was then added into 40 μ L of each inhibition/reversal assay mixture, thoroughly vortexed and transferred into an appropriate cuvette. The fluorescence of amyloid-bound ThT was measured by

fluorescence spectroscopy using a Cary Eclipse fluorescence spectrophotometer; bound ThT exhibits a new excitation maxima at 450 nm and an enhanced emission maxima at 482 nm.³³

OAF microscopy experiments

Samples were imaged on a custom built OAF²⁵ system. The OAF system permits high signal to noise imaging with greater sample depth penetration. Fluorescence imaging was achieved through a 500LP dichroic and clean up filter (Chroma, VT) before entering an Optosplit III triple color image splitter (Cairn, UK), and was detected using an EMCCD camera (Andor DU897, Andor, UK). The emission wavelength range used was 500–565 nm, therefore, both the excitation and emission wavelengths were off peak for ThT, however, the image quality was excellent. All samples were prestained with 10 μ M ThT, pipetted onto a clean glass slide, air dried and then imaged in 10 mM potassium phosphate buffer, pH 7.4, supplemented with 100 mM DTT to minimize photobleaching. For consistency and cross-correlation, the same samples were used for inhibition/reversal imaging as those in ThT and CD experiments. In addition, all OAF experiments were performed blind to prevent bias toward any one sample.

AFM experiments

Samples were imaged in noncontact mode using a XE-120 Atomic Force Microscope (Park Systems, South Korea). NSC 15 silicon nitride cantilevers with a spring constant of 40 N/m were used for imaging at a scan rate of 1.0 Hz and a resolution of 256×256 pixels. All images were taken at room temperature. The AFM data were taken from inhibitors alone as well as inhibition and reversal experiments, using *cys*₁₅₂₁, *cys*_{2935a}, and the *cys*₁₅₂₁ + *cys*_{2935a} mixture. 5 μ L sample was placed on freshly cleaved mica (thickness 0.3 mm). Following adsorption of the protein aggregates (2 min), the mica was washed with 5 mL of double distilled water. Excess water was removed and the samples were dried using a stream of nitrogen gas. Samples were immediately analyzed by AFM.

3-(4,5 Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide cell-toxicity assay

MTT experiments were undertaken using Rat pheochromocytoma (PC12) cells to assess the effect of the toxicity of A β ₄₂. PC12 cells are known to be particularly sensitive and their use in this assay is well established.³⁴ The MTT Vybrant® MTT Cell Proliferation Assay Kit (Invitrogen) was used to measure the conversion of the water soluble MTT dye to formazan, which is then solubilized, and the concentration determined by a color change monitored via absorbance measurement at 570 nm. The change in absorbance can then be converted to a percentage MTT reduction which can be used as an indicator of the

PC12 cell health in the assay. The assay was performed with 10 μ M A β ₄₂ and varying molar ratios of peptide corresponding to 1:0.1 (0.1 μ M), 1:1 (10 μ M), 1:4 (40 μ M), 1:10 (100 μ M). PC12 cells were maintained in RPMI 1640 + 2 mM Glutamine medium mixed with 10% Horse Serum, 5% Fetal Bovine Serum, supplemented with a 20 mg/mL Gentamicin. Cells were transferred to a sterile 96-well plate with 30,000 cells per well and experiments performed in triplicate. Briefly, different concentrations of peptides were screened in the presence of 10 μ M A β ₄₂. The required volume from peptide and target stock solutions was freeze-dried overnight. The freeze-dried peptide and A β ₄₂ target were resuspended in 100% dimethyl sulfoxide (DMSO), each at 100 \times stock concentration (i.e., 1, 2, 4, or 10 mM). For example, for the molar ratio 1:1 a total of 5 μ L from each of the resuspended inhibitor/DMSO and target/DMSO was mixed in a well of a 96-well preparation plate, thus giving 10 μ L of 1:1 mM inhibitor/target concentration ratio in 100% DMSO. A total of 90 μ L of RPMI media was added to the 10 μ L inhibitor/target mixture (100:100 μ M inhibitor/target ratio in 10% DMSO). A total of 10 μ L of the 50:50 μ M inhibitor/target mixture in 10% DMSO was then dispensed into 90 μ L of media/PC12 cells, at final peptide inhibitor and target concentrations of 10 μ M. These were incubated for 24 h at 37°C, 5% CO₂, prior to the addition of the MTT dye. A total of 10 μ L of the dye was added to each well and incubated for a further 4 h at 37°C, 5% CO₂. A total of 100 μ L of the DMSO (stop/solubilization solution) was then added to each well and was allowed to stand for 10 min. The absorbance was measured at 570 nm using a 96-well Versamax tunable microplate reader. Assay values for incubation with buffer were taken as 100% and inhibition of cell function by incubation with buffer containing A β ₄₂ alone taken as 0%. Raw data were then scaled as follows: (raw data point – A β ₄₂ mean)/(buffer mean – A β ₄₂ mean) \times 100. The scaled mean for each data set was then plotted with the error given by the standard deviation.

Acknowledgments

The authors acknowledge the use made of the Park Systems XE-120 Atomic Force Microscope which was on loan from the EPSRC (Engineering and Physical Sciences Research Council) Engineering Instrument Pool. The authors wish to thank Dr Miao Yu and Dr Victoria Allen-Baume for excellent technical support throughout the project.

References

1. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL (1998) Diffusible, nonfibrillar ligands derived from A β _{1–42} are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA* 95:6448–6453.

2. Klein WL, Krafft GA, Finch CE (2001) Targeting small Abeta oligomers: the solution to an Alzheimer's disease conundrum? *Trends Neurosci* 24:219–224.
3. Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8:101–112.
4. Walsh DM, Selkoe DJ (2007) A beta oligomers—a decade of discovery. *J Neurochem* 101:1172–1184.
5. Comery TA, Martone RL, Aschmies S, Atchison KP, Diamantidis G, Gong X, Zhou H, Kreft AF, Pangalos MN, Sonnenberg-Reines J, Jacobsen JS, Marquis KL (2005) Acute gamma-secretase inhibition improves contextual fear conditioning in the Tg2576 mouse model of Alzheimer's disease. *J Neurosci* 25:8898–8902.
6. O'Nuallain B, Freir DB, Nicoll AJ, Risse E, Ferguson N, Herron CE, Collinge J, Walsh DM (2010) Amyloid beta-protein dimers rapidly form stable synaptotoxic protofibrils. *J Neurosci* 30:14411–14419.
7. Sandberg A, Luheshi LM, Sollvander S, Pereira de Barros T, Macao B, Knowles TP, Biverstal H, Lendel C, Ekholm-Pettersson F, Dubnovitsky A, Lannfelt L, Dobson CM, Hard T (2010) Stabilization of neurotoxic Alzheimer amyloid-beta oligomers by protein engineering. *Proc Natl Acad Sci USA* 107:15595–15600.
8. Petkova AT, Ishii Y, Balbach JJ, Antzutkin ON, Leapman RD, Delaglio F, Tycko R (2002) A structural model for Alzheimer's beta-amyloid fibrils based on experimental constraints from solid state NMR. *Proc Natl Acad Sci USA* 99:16742–16747.
9. Hard T, Lendel C (2012) Inhibition of amyloid formation. *J Mol Biol* 421:441–465.
10. Acerra N, Kad NM, Mason JM (2013) Combining Intracellular selection with protein-fragment complementation to derive Aβ interacting peptides. *Protein Eng Des Sel* 26:463–470.
11. Acerra N, Kad NM, Griffith DA, Ott S, Crowther DC, Mason JM (2014) Retro-inversal of intracellular selected ABeta interacting peptides: implications for a novel Alzheimer's disease treatment. *Biochemistry* 53:2101–2111.
12. Sharpe S, Yau WM, Tycko R (2005) Expression and purification of a recombinant peptide from the Alzheimer's beta-amyloid protein for solid-state NMR. *Protein Expr Purif* 42:200–210.
13. Soscia SJ, Kirby JE, Washicosky KJ, Tucker SM, Ingelsson M, Hyman B, Burton MA, Goldstein LE, Duong S, Tanzi RE, Moir RD (2010) The Alzheimer's disease-associated amyloid beta-protein is an antimicrobial peptide. *PLoS One* 5:e9505.
14. Mason JM, Schmitz MA, Muller KM, Arndt KM (2006) Semirational design of Jun-Fos coiled coils with increased affinity: Universal implications for leucine zipper prediction and design. *Proc Natl Acad Sci USA* 103:8989–8994.
15. Remy I, Campbell-Valois FX, Michnick SW (2007) Detection of protein-protein interactions using a simple survival protein-fragment complementation assay based on the enzyme dihydrofolate reductase. *Nat Protoc* 2:2120–2125.
16. Pike CJ, Walencewicz-Wasserman AJ, Kosmoski J, Cribbs DH, Glabe CG, Cotman CW (1995) Structure-activity analyses of beta-amyloid peptides: contributions of the beta 25–35 region to aggregation and neurotoxicity. *J Neurochem* 64:253–265.
17. Hughes E, Burke RM, Doig AJ (2000) Inhibition of toxicity in the beta-amyloid peptide fragment beta-(25–35) using N-methylated derivatives: a general strategy to prevent amyloid formation. *J Biol Chem* 275:25109–25115.
18. Hung LW, Ciccotosto GD, Giannakis E, Tew DJ, Perez K, Masters CL, Cappai R, Wade JD, Barnham KJ (2008) Amyloid-beta peptide (Abeta) neurotoxicity is modulated by the rate of peptide aggregation: Abeta dimers and trimers correlate with neurotoxicity. *J Neurosci* 28:11950–11958.
19. Tjernberg LO, Naslund J, Lindqvist F, Johansson J, Karlstrom AR, Thyberg J, Terenius L, Nordstedt C (1996) Arrest of beta-amyloid fibril formation by a pentapeptide ligand. *J Biol Chem* 271:8545–8548.
20. Lowe TL, Strzelec A, Kiessling LL, Murphy RM (2001) Structure-function relationships for inhibitors of beta-amyloid toxicity containing the recognition sequence KLVFF. *Biochemistry* 40:7882–7889.
21. Zagorski MG, Yang J, Shao H, Ma K, Zeng H, Hong A (1999) Methodological and chemical factors affecting amyloid beta peptide amyloidogenicity. *Methods Enzymol* 309:189–204.
22. Soto C, Kindy MS, Baumann M, Frangione B (1996) Inhibition of Alzheimer's amyloidosis by peptides that prevent beta-sheet conformation. *Biochem Biophys Res Commun* 226:672–680.
23. Soto C, Sigurdsson EM, Morelli L, Kumar RA, Castano EM, Frangione B (1998) Beta-sheet breaker peptides inhibit fibrillogenesis in a rat brain model of amyloidosis: implications for Alzheimer's therapy. *Nat Med* 4: 822–826.
24. Kokkoni N, Stott K, Amijee H, Mason JM, Doig AJ (2006) N-Methylated peptide inhibitors of beta-amyloid aggregation and toxicity. Optimization of the inhibitor structure. *Biochemistry* 45:9906–9918.
25. Kad NM, Wang H, Kennedy GG, Warshaw DM, Van Houten B (2010) Collaborative dynamic DNA scanning by nucleotide excision repair proteins investigated by single-molecule imaging of quantum-dot-labeled proteins. *Mol Cell* 37:702–713.
26. Levy R, Weiss R, Chen G, Iverson BL, Georgiou G (2001) Production of correctly folded Fab antibody fragment in the cytoplasm of *Escherichia coli* trxB gor mutants via the coexpression of molecular chaperones. *Protein Expr Purif* 23:338–347.
27. Mason JM, Gibbs N, Sessions RB, Clarke AR (2002) The influence of intramolecular bridges on the dynamics of a protein folding reaction. *Biochemistry* 41:12093–12099.
28. Maurer-Stroh S, Debulpaep M, Kuemmerer N, Lopez de la Paz M, Martins IC, Reumers J, Morris KL, Copland A, Serpell L, Serrano L, Schymkowitz JW, Rousseau F (2010) Exploring the sequence determinants of amyloid structure using position-specific scoring matrices. *Nat Methods* 7:237–242.
29. Frousios KK, Iconomidou VA, Karletidi CM, Hamodrakas SJ (2009) Amyloidogenic determinants are usually not buried. *BMC Struct Biol* 9:44.
30. Trovato A, Seno F, Tosatto SC (2007) The PASTA server for protein aggregation prediction. *Protein Eng Des Sel* 20:521–523.
31. Tartaglia GG, Vendruscolo M (2008) The Zyggregator method for predicting protein aggregation propensities. *Chem Soc Rev* 37:1395–1401.
32. Fernandez-Escamilla AM, Rousseau F, Schymkowitz J, Serrano L (2004) Prediction of sequence-dependent and mutational effects on the aggregation of peptides and proteins. *Nat Biotechnol* 22:1302–1306.
33. LeVine H, III (1993) Thioflavine T interaction with synthetic Alzheimer's disease beta-amyloid peptides: detection of amyloid aggregation in solution. *Protein Sci* 2:404–410.
34. Shearman MS, Ragan CI, Iversen LL (1994) Inhibition of PC12 cell redox activity is a specific, early indicator of the mechanism of beta-amyloid-mediated cell death. *Proc Natl Acad Sci USA* 91:1470–1474.